

MAGNITUDE OF CLINICAL PROBLEM AND CLINICAL PROFILE OF POISONING IN PEDIATRICS

Bhavya E & Chakrapani B *

Department of Pharmacy Practice, School of Pharmaceutical Sciences, Vel's Institute of Science Technology and Advanced Studies (VISTAS), Chennai

** Address for correspondence: Mr. B. Chakrapani, Research Scholar, Department of Pharmacy Practice, School of Pharmaceutical Sciences, Vel's Institute of Science Technology and Advanced Studies (VISTAS), Chennai, India. E-mail ID: chakrispecial@gmail.com, Mob: 9618279507*

ABSTRACT

Background: *Children are inquisitive and explore their world with all their senses. Poisoning patterns change according to age group, the nature and dose of the poison, type of exposure. This study is aimed at understanding the incidence, aetiology and manner of poisoning, outcome among children less than 18 years admitted to Paediatric department.*

Methods: *Prospectively, 38 poisoning cases admitted to Paediatric intensive care, from August 2021 to May 2022 were included in this study. The age and sex of the patient, poisoning agent, manner of poisoning, duration of hospital stay and outcome of all the patients were recorded and analyzed.*

Results: *Out of 38, 22 (58%) are males, 16 (42%) are females. 13 (34%) cases are <5 years of age, 2(5%) cases are between 5-10 year, 23 (61%) cases are above 10 years. Among all, 13 (34%) cases of drugs consumption, among these 11 cases were above 10 years representing purposeful consumption. 5 (13%) cases of kerosene consumption. Out 4 (11%) cases of Organophosphorous poison, 3 cases are above 10 years representing purposeful consumption. 4 (10%) cases of acid consumption. 4 (11%) cases of aluminium phosphide poisoning. 3 (8%) cases of Good night oil consumption. 2 (5%) cases of eucalyptus oil consumption. 3(8%) cases of camphor poisoning. 2 (5%) cases died due to aluminium phosphide poisoning. Duration of hospital stay varied from 1 day to 6 days.*

Conclusions: *Accidental poisoning is common in children below 5 years of age. Suicidal poison incidence is increasing among teenagers. Accidental poisoning can be reduced by simple measures like parental education, replacing the poisoning agent with one of lower toxicity, legislation regarding the child resistant packaging of necessary poisons.*

Keywords: *Camphor, Drugs, Eucalyptus oil, Poisons*

INTRODUCTION

The cause and poisoning type vary in different parts of the world and within the country also depending upon factors such as education, demography, socioeconomic factors, customs and local belief.¹ The mortality and morbidity due to poisoning is preventable in children. Poisoning is the fourth leading cause of unintentional injury following road traffic accident, burns and drowning.² Approximately 2 million people each year under 6 years come to emergency department with history of poisoning.³ Although Poisoning never accounted for a large number of accidental deaths, the rate of decrease in deaths due to poisoning have not been as drastic as infectious diseases, hence have assumed more prominence now.⁴⁻⁶ Pattern of Poisoning varies according to age, the nature and dose of the poison and type of exposure.⁷ Among the accidental poisoning, even though there is no significant decrease in the number, the pattern of poisoning has shown a change with new hazards constantly appearing due to introduction of newer drugs and chemicals for domestic use, farming.⁸ The most common agents involved in poisoning are over-the-counter (OTC) medications, prescription medications, household products, pesticides, kerosene, poisonous plants and animal or insect bites.⁹ In acute poisoning, sudden onset of symptoms appear after the ingestion, inhalation or after coming in contact with poisonous substance. In chronic poisoning, symptoms develop gradually over a period of time and there is complete disappearance of symptoms on the removal of patients from exposure. Acute poisoning in childhood is an important cause of morbidity and mortality and can be effectively controlled by preventive measures.¹⁰ The most common cause of acute medical illness in many countries has been exposure to toxic substances.¹¹⁻¹³ General epidemiological data should be used to assist emergency department on proper management of poisoning cases especially for effective preventive and therapeutic approaches.¹⁴⁻¹⁵

METHODS

This is a Prospective observational study conducted in Department of Paediatrics, between August August 2021 to May 2022. All children and adolescent who were admitted to pediatric intensive care unit with history of poisoning between August 2021 to May 2022 were included in this study. Total 52 cases came to emergency with history of poisoning during this period. We included all children below 18 years who came with history of poisoning and who had signs and symptoms of poisoning at admission. Those cases with history of contact poisons, food poisoning, animal bite, insect bite, chronic poisoning and those having no signs and symptoms of poisoning were excluded from the study. Out of 52 cases only 38 cases met our inclusion criteria. Study was conducted after obtaining verbal and written consent from the parent or guardian. These cases admitted to Pediatric intensive care after initial assessment. In each case detailed history was taken-time when poison was ingested or come in contact with it, substance taken, its dose, cause for poisoning, manner of poisoning, any intervention done before reaching our hospital and time brought to casualty was noted. Detailed general physical examination and systemic examination done. Data regarding age of the patient, sex, etiological agent, nature of poisoning (accidental or intentional), clinical presentation, duration of hospital stay and outcome were collected and filled in a predesigned Performa. The data collected were then transferred into excel format

and analysed as percentage.

RESULTS

Total 513 cases admitted to PICU during our study period. Out of these 52 cases came with history of poisoning. Out of which 38 cases met our inclusion criteria. The incidence of poisoning in our study found to be 1.29%. Coming to sex distribution, 22 cases were males accounting for 58%, 16 cases were females accounting for 42% as shown in Figure 1.




Figure 1: Distribution of sex

Duration of hospital stay ranged from 0-6 days. 13 cases (34.2%) stayed in hospital ranging between 0-2 days. 22 cases (57.9%) stayed in hospital for 2-4 days. Only 3 cases (7.9%) stayed between 4-6 days.

Table 1: Duration of hospital stay

Duration	No. of cases	Percentage	P-Value
0-2 days	13	34.2%	0.00
2-4 days	22	57.9%	0.00
4-6 days	03	7.9%	0.82

In this study time taken to reach hospital from the time they consumed poison was also calculated. Majority of the cases reached within 1 to 4 hour accounting for 47.3%, 12 cases reached within 1 hour of consumption accounting for 31.5%. 8 cases reached after 4 hours accounting for 21.05%

Table 2: Time since poison consumption to arrival at casualty

Time	No. of cases	P-Value
Less than 1 hour	12(31.5%)	0.00
Between 1 to 4 hour	18(47.3%)	0.00
More than 4 hour	8(21.05%)	0.08

Significant $p \leq 0.01$

Time taken to reach emergency since poison consumption is very important. Because the early the intervention better is the outcome. Those cases which presented early can be given gastric lavage to prevent further absorption. However, 2 cases of aluminum phosphide poison died even though they came to emergency within 1 hour of consumption. Out of 38 cases of poisoning, 2 cases (5.2%) died both due to aluminum phosphide poisoning. In both these cases we did gastric lavage immediately but these cases died within half an hour of presentation. Duration of hospital stay ranged from 0-6 days. 13 cases (34.2%) stayed in hospital ranging between 0-2 days. 22 cases (57.9%) stayed in hospital for 2-4 days. Only 3 cases (7.9%) stayed between 4-6 days. Among the cases admitted, drug ingestion accounted for maximum number of cases 34.21%. Among the drug ingestion 4 cases of anti-epileptics ingestion, 3 cases of anti-histamines, 2 cases of anti-hypertensives, 2 cases of thyroid drugs and 2 cases of antipsychotic and anti-depressants each. Among 8 (21%) insecticides and pesticide cases, 4 were organophosphates and 4 were aluminium phosphide poisoning. Among the others, kerosene consumption were 5 cases, 4 cases were cleaning agent consumption, 3 cases of good night oil consumption, 2 cases each of camphor and eucalyptus oil consumption and 1 cannabis poisoning. We can observe the easily available household things like kerosene and good night oil are the poisoning agents involved in children below 5 years representing accidental poisoning. On the other hand, drugs as etiological agent more frequently found in teenagers, representing suicidal poison.

Most common symptoms at the time of presentation, vomiting being the most common symptom noted in 28 cases accounting for 73.7%, followed by altered sensorium present in about 47.4% of cases. Restlessness was present in 42.1% of cases. Poison odor in 36.8% of cases. Headache and Pain abdomen in 13.2% of cases each. Respiratory distress at the time of presentation was present in 7.9% of cases. Convulsion and giddiness were present in 5.3% of cases.

Table 3: Etiological agent involved

Poisoning agent	Type of poison	No. of cases	P-Value
Insecticides and pesticides	organophosphates	04	0.11
	Aluminium phosphide	04	0.18
Drugs	Anti epileptics	04	0.18
	Anti histamines	03	0.16
	Anti hypertensives	02	0.22
	Thyroid drugs	02	0.22
	Antipsychotics and anti depressants	02	0.23
Cleaning agent	Acid	04	0.19
Kerosene		05	0.16
Good night oil		03	0.18
Eucalyptus oil		02	0.12
Camphor		02	0.11
Cannabis		01	0.10

Significant $p \leq 0.01$

Table 4: Presenting symptoms

Symptom	No. of cases	p- value
Vomiting	28	0.00
Altered sensorium/Unconsciousness	18	0.00
Restlessness	16	0.00
Odour	14	0.00
Headache	05	0.18
Pain abdomen	05	0.22
Respiratory distress	03	0.36
Convulsions	02	0.59
Giddiness	02	0.54

Significant $p \leq 0.01$

DISCUSSION

Poisoning in children is important cause of morbidity and mortality in developing countries like India. Our study aimed at describing magnitude of clinical problem and clinical profile of poisoning cases. In this study, the incidence of poisoning found to be 1.29%. Similar incidence has been reported in other studies.¹⁶ With respect to sex preponderance, incidence was more common in males in less than 5 years of age group, whereas female predominance was seen in adolescent age group; with overall female predominance, while other studies showed overall incidence to be more common in males.¹⁷ Majority of the children are aged above 10 years (60.5%) followed by less than 5 years (34.2%). Similar results are observed in study done by Sharma et al.²⁵ Other studies reported the most common age group being 1-6 years.¹⁸ Study done by Singh M et al reported maximum number of cases in children aged above 5 years.¹⁹

Most of the cases (47.3%) arrived to casualty between 1 and 4 hours of poison consumption. A study done from south East Asia also reported most cases presented between 1 and 6 hour of poisoning.²⁰ A study done by Ahmed et al reported most of the cases (80%) arrived to casualty within 1 hour of consumption.²¹ The earlier the arrival to emergency department better will be the outcome. This is because we can intervene earlier. Duration of hospital stay varied from 0 to 6 days. Most of the cases stayed in ICU ranging 2 to 4 days (57.9%). Almost similar results were reported in another study where most cases stayed ranging 1 to 3 days (55.03%).^{22,23} Coming to clinical presentation, there was wide variety of clinical presentation according to type of poison and age group. However most common presenting complaint being vomiting (73.7%), altered sensorium (47.4%), restlessness (42.1%), odour from the mouth (36.8%), Headache (13.2%). Almost similar presenting complaints observed in study done by Sharma J et al.²⁴ Other studies also reported the most common presenting complaint being gastrointestinal followed by central nervous system.

The other studies done in past showed kerosene is the most common poisoning agent accounting for 25%- 50%.²⁴ But in our study kerosene is the 3rd most common poisoning agent. This probably due use of gas instead kerosene for cooking. A study done by Vasanthan et al reported kerosene being the most common etiological agent followed by drugs.²⁵ The

other studies reported drugs followed by kerosene are the most common poisoning agent in their study.²⁶ Out of 38 cases, 2 cases died. Both cases are of alluminium phosphide poisoning.

CONCLUSION

Suicidal poisoning has been shown to be more predominant in adolescents and teenagers especially girls, hence need for a guidance and counselor at every school to help child deal with stressful situations, and to guide parents as well. In either case, the most important step is to bring child to the emergency room at the earliest this has been proven reduce the morbidity and mortality significantly.

REFERENCES

1. Narayan Reddy KS. The essential of Forensic Medicine and Toxicology, 27th edition, Medical Book Company. 2008;95:440-456.
2. Hyder AA, Wali S, Fishman S, Schenk E. The burden of unintentional injuries among the under-five population in South Asia. *Acta Paediatrica*. 2008;97:267-75.
3. American Academy of Pediatrics. Poison Prevention and Treatment Tips National Poison Prevention Week, March 19-25;2017.
4. Parikh CK. Toxicology. In: Parikh CK (ed). Parikh's text book of medical jurisprudence and toxicology. 5th Edn. Bombay. CBS publisher; 1990:660- 690.
5. Modi NJ. Toxicology. In: Modi NJ (ed). Modi's Textbook of medical jurisprudence and toxicology. 20th Edn. Bombay. NM Tripathi Pvt Ltd.; 1980:455-470.
6. Reddy NKS. Toxicology. In: Narayan Reddy KS (ed). The synopsis of forensic medicine and toxicology. 14th Edn. Hyderabad. Medical book company; 2000:221-293.
7. WHO-UNICEF. Children and poisoning: world report on child injury prevention. World Health Organization. 2008.
8. Singh S, Singhi S, Sood NK, Kumar L, Walia BNS. Changing pattern of childhood poisoning (1970- 1989): Experience of a large North Indian hospital. *Indian Pediatr*. 1995;32:331-6.
9. Lam LT. Childhood and adolescence poisoning in NSW, Australia: an analysis of age, sex, geographic, and poison types. *Inj Prev*. 2003;9:338-42.
10. Litovitz TL, Klein-Schwartz W, Rodgers GC. Annual report of American association of poison control centers toxic exposure surveillance system. *Am J Emerg Med* 2002; 20: 391-452.
11. The global burden of disease: A comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020. Murray, CJ and Lopez, AD, eds. Harvard School of Public Health (on behalf of the World Health Organization and The World Bank); 1996:120-122.
12. Ferguson RW, Mickalide AD. An In-Depth Look at Keeping Young Children Safe Around Medicine. Washington, DC: Safe Kids Worldwide, March 2013.
13. Juris E. Personal communication. Washington, DC: American Association of Poison Control Centers;2006.
14. Consumer Product Safety Commission. CPSC warns that 9 out of 10 unintentional child

poisonings occur in the home. News from CPSC, March 18, 2009. U.S. Consumer Product Safety Commission Website. Available from:

<http://www.cpsc.gov/cpscpub/prerel/prhtml09/09159.html>. Accessed: November 3, 2014.

15. Dutta AK, Seth A, Goyal PK, Agarwal V, Mittal SK, Sharma R et al. Poisoning in children, Indian scenario. *Indian J Pediatr.* 1998;65:365-70.
16. Krishankumar P, Geeta MG, Gopalan AV. Deliberate Self-Poisoning in Children. *Indian Pediatr.* 2005;42:582-6.
17. Budhathoki S, Poudel P, Bhatta NK, Dutta AK, Shah GS, Bhurtyel KK, et al. Clinical profile and outcome of poisoning and intoxication in children: A hospital based study. *Nepal Med Coll J.* 2009;11(3):170-5.
18. Shoter AM. Kerosene poisoning in childhood: a 6-y prospective study at the Princess Rahmat Teaching Hospital. *Neuro Endocrinol Lett.* 2005;26:835-38.
19. Rashid Mamunur AKM, Sultana R, Nazmul Ahasan HAM, Rasul CH, Seasonal variation of acute childhood Poisoning, *Pak J Med Sci.* 2007;23(3):443-5.
20. Desalew M, Aklilu A, Amanuel A, Addisu M, Ethiopia T. Pattern of acute adult poisoning at Tikur Anbessa specialized teaching hospital, A retrospective study, Ethiopia. *Hum Exp Toxicol.* 2011;30:523-7.
21. Ahmed A, Aljamal A, Ibrahim M. Poisoning emergency visits among children: A 3-year retrospective study in Qatar. *BMC Pediatr.* 2015;15,104:1-7.
22. Bhat NK, Dhar M, Ahmad S, Chandar V. Profile of poisoning in children and adolescents at a North Indian tertiary care centre. *JIACM.* 2011;13:37-42.
23. Sitaraman S, Sharma U, Saxena S. Accidental Poisoning in children. *Indian Paediatrics.* 1985;22:757-60.
24. Sharma J, Kaushal RK. Profile of poisoning in children. *Pediatric Oncall.* 2014;11:40-2.
25. Vasanthan M, James S, Shuba S, Abhinaya J. Clinical profile and outcome of poisoning in children admitted to a tertiary referral center in South India. *Indian J Child Health.* 2015;2(4):187- 91.
26. Akhtar S, Rani GR, Al-Anezi FA. Risk factors in acute poisoning in children-A retrospective study. *Kuwait Med J.* 2006;38(1):33-6.