

Climate Change and Pollinator Dynamics: Integrating Social Media Insights and Ecological Data for Conservation Strategies

Pooja Hadimane¹, Ashoka K², Basavaraja A S³

¹*Department of CSE, Research Scholar, Bapuji Institute of Engineering and Technology
(Visvesvaraya Technological University, Belagavi), Davangere, India*

²*Department of ISE, Professor, Bapuji Institute of Engineering and Technology
(Visvesvaraya Technological University, Belagavi), Davangere, India*

³*Department of ECE, Lecturer, Gov. Polytechnic, Harihar, India*

Abstract

Pollination is an essential ecosystem service intricately linked to biodiversity, ecosystem health, and agricultural systems. The need to understand the effect of climate change on pollination processes has never been greater, given that a significant portion of global crop production is dependent on biotic pollination. This survey paper examines the multifaceted challenges that climate change poses to pollination dynamics across various ecosystems. By synthesizing existing literature to highlight how alterations in temperature and precipitation patterns have led to a phenological mismatch between pollinators and plants, potentially disrupting established trophic relationships and ecosystem functions. Our review reveals that insect-pollinated plants, particularly those that bloom early in the season, exhibit a heightened sensitivity to climate-induced phenological shifts. Moreover, exploring how the altered life cycles of pollinators, struggling to synchronize with the new flowering schedules, may precipitate declines in pollination services. Our findings underscore the critical need for conservation strategies that address climate adaptation for pollinators, focusing on enhancing landscape connectivity and heterogeneity. By bridging diverse studies ranging from the application of social media data in ecological research to advanced predictive models for pollination services, the main aim is to foster a deeper understanding of the consequences of climate change on pollination.

Keywords: Pollination, Ecosystem health, Pollination dynamics, Phenological mismatch, Climate Change

1 INTRODUCTION

Pollination is an ecological function that is extremely valued economically and directly tied to human well-being since it influences agricultural productivity and food security. Pollinators affect the world's food supply; pollinator-dependent crops make up around 35% of total agricultural production by volume. Estimates provided state that biotic pollination is necessary for the production of fruits and seeds in 87 out of the 115 major crops farmed worldwide [1]. Additionally, it is shown that animal pollination services are necessary for about 90% of wild flowering plant species.

Furthermore, animal pollination is essential to over 75% of the world's primary food crop varieties for either their quantity or quality, or both. Recognizing the importance of pollination for the survival of contemporary people and for guaranteeing food security is crucial. The study highlighted the importance of pollinator and pollination monitoring on a global scale. The assessment of pollinators, pollination, and food production by IPBES also brought attention to data gaps and emphasized how important it is to monitor these areas [2]. Furthermore, the assessment revealed indications of a decline in natural pollinators globally. The production of genetically modified (GM) crops, the introduction of invasive species, the widespread use of pesticides, the expansion and intensification of agricultural practices, the presence of parasites, and many other factors all contribute to the decline of pollinators [3]. Moreover, habitat loss and fragmentation brought on by human activities, such as greater intensification, also contribute to the decline in pollinators. Accordingly, pollination is one common ecological service and function that is directly impacted by the continuous evolution of landscapes influenced by people [4]. In human-altered habitats, pollinators and flowering plants—especially those with specific reproductive traits—are frequently driven to local extinction or drastically reduced. Additionally, amid highly fragmented forest remnant landscapes, geographically isolated populations impede pollen exchange across populations, hence decreasing the success rate of plant reproduction [5]. Agriculture has a favorable effect on pollinator abundance, but it is also the primary cause of the decline in pollinator populations. This loss is attributed to shifts in land use, the use of agrochemicals, and the adoption of conventional or automated large-scale agricultural operations [6]. Figure 1 shows the pollination process.




Figure 1 pollination process

Types of Pollination

- **Wind-Pollinated:** If the flower is wind-pollinated, the flowchart directs the process to where the wind carries the pollen to another flower, resulting in pollination.
- **Animal-Pollinated:** If the flower is not wind-pollinated, the next question asks if it is animal-pollinated. If it is, an animal (most likely an insect) visits the flower for nectar. During the visit, pollen sticks to the animal. When the animal visits another flower, the pollen is transferred to this second flower, achieving pollination.
- **Self-Pollination:** If the flower is not animal-pollinated (and by extension, not wind-pollinated), self-pollination occurs within the same flower, leading to pollination.

Several studies [7-8] have indicated that one of the most important disturbance factors induced by humans that influence ecosystems is climate change. Regarding the possible effects of climate change on interactions between plants and pollinators, a number of hypotheses have been put out. The physiological vulnerability of organisms to changes in different environmental circumstances determines the impacts of climate change. Recent study has shown that the phenomena of global warming significantly affects the phenology and distribution of different plants and animals. The finding that spring events occur 2.3 days' sooner on average worldwide and that species range moves towards the poles at a pace of 6.1 km per decade provide evidence in favor of this argument. As temperatures rise, studies have shown alterations in trophic relationships and energy fluxes in plant-herbivore and predator-prey interactions. Although there hasn't been a commensurate adjustment in the date of plant blossoming found a substantial temporal shift in pollinator emergence in alpine settings in recent years [9]. In general, plants that get insect pollination react to heat more strongly than plants that receive wind pollination.

Studies suggest that early-blooming species of these plants are the most temperature-sensitive. This implies that temperature variations have a significant impact on the phenologies of these specific species. Strong evidence has been gathered over the past few decades that many plant communities in Europe have seen longer growing seasons. Climate change is having an effect on plants' phenological responses, which might make it difficult for pollinators to synchronize their life cycles with the altered timing of pollination. The study has demonstrated a continuous correlation between the average temperature during the month of flowering or the months before to it and the commencement of blooming. Many studies have demonstrated that the link between the commencement of flowering and rising temperatures is, for the most part, linear [10]. This discovery may affect how pollinators and plants interact, as well as how other creatures interact with plants. Furthermore, a number of other variables, including photoperiodicity, precipitation, soil humidity, and snowmelt, might affect the onset of flowering. Moreover, specific signal combinations may potentially function as possible triggers for blooming start. Because species vary in their mobility and capacity to adapt to changing climatic conditions, it is anticipated that the composition of plant and pollinator assemblages would shift in many different locales. Tropical species are found to live in habitats that are either at or very near their ideal temperature range. Certain species may be forced to relocate to cooler areas or run the risk of going extinct if the temperature rises further. Because pollinators react behaviourally to avoid extreme temperatures, future global warming may have a substantial impact on pollination services. Since most pollinators are insects, temperature probably has a considerable impact on their life cycle, growth, and patterns of activity [11]. Their tiny size and poikilothermic nature are the main causes of this.

A research on butterflies found that temperature and the date of the butterflies' first emergence were significantly correlated. Climate cues like temperature and water availability affect when pollination occurs. Insect pollinators' foraging behavior is influenced by environmental conditions. These variables include when foraging begins and ends, how many plants and flowers pollinators visit, and how many pollinators there are in each square meter. They are directly impacted by the aforementioned variables, which include rising temperatures, erratic rainfall patterns, and other environmental variables. The estimated that over 80% of pollination tasks are carried out by insects. Climate change so has a detrimental effect on agricultural yield and might be a serious danger to the world's food security.

The research is driven by the critical ecological and economic importance of pollinators in global agriculture and food security, recognizing that a significant portion of crop production relies on biotic pollination. With evidence of pollinator decline due to factors such as pesticide use, habitat loss, and climate change, there is a pressing need to understand and mitigate these impacts. Climate change, in particular, threatens to disrupt plant-pollinator interactions by altering phenology and geographical distributions, potentially leading to mismatches in pollination timing and jeopardizing crop yields. This research seeks to fill the data gap in pollinator monitoring, particularly in under-studied regions, and aims to inform conservation strategies that currently lack comprehensive climate change mitigation measures. The urgency of addressing this issue is compounded by the implications for food security, as the majority of pollination is carried out by insects, which are especially sensitive to environmental changes.

- The research provides an in-depth analysis of the multifaceted causes behind pollinator decline, specifically highlighting the roles of pesticide use, habitat fragmentation, and climate change, thereby contributing valuable insights into the conservation and management of pollinator populations essential for food security and biodiversity.
- By identifying the urgent need for comprehensive monitoring and research, particularly in regions where data is scarce, the study offers a roadmap for future research directions, stressing the importance of global efforts in tracking and understanding pollinator behaviors and populations in the face of environmental changes.
- The study critically evaluates current conservation strategies, revealing a significant oversight regarding climate change mitigation efforts, and underscores the need for integrating climate adaptation initiatives into pollinator conservation plans to ensure the sustainability of these vital species in the Northern Hemisphere.

2 RELATED WORK

Globally, pollinator biodiversity is declining at a rate never seen before, as of right now, the main reasons for the decline are up for debate. Although the primary cause of the predicted future reduction in biodiversity is expected to be climate change, the possible beneficial effect of present warming on pollinator assemblages at higher latitudes is being discussed. Pollinator conservation initiatives sometimes lack climate adaption measures. When pollinator health is greatly impacted by climate change, there is a potential risk. The literature that has been written about variety at the genetic, species, and community levels has been thoroughly examined [12]. Our findings suggest that as a result of global warming, pollinator assemblages at all pollinator biodiversity levels will become homogenized. Their resilience to subsequent stochastic occurrences is weakened by this homogeneity. The effect is mostly noted on aspects of biodiversity that are frequently not measured, such genetic diversity, β -diversity, and species evenness.

Furthermore, certain aspects of climate change—like variations in winter temperatures, an extension of the vegetative season, and an increase in the frequency of extreme weather events—are frequently understudied despite having a particularly negative effect on pollinators. It is thought that the unfavourable impacts of global warming on pollinator biodiversity are made worse in Europe and the US by the prevalence of uniform and fragmented environments [13]. The main causes of this are the scarcity of range-shifting and microclimatic buffering possibilities. Improving landscape connectedness and variety at different geographic scales need to be the main goal of conservation initiatives.

Study [7] emphasizes the use of social media data—Flickr images specifically—for the analysis of leisure travel in the Tahoe Central Sierra Initiative project region in California. The application of machine learning techniques facilitates the recognition of noteworthy natural occurrences and the forecasting of prospective leisure prospects. This method emphasizes how important it is to use user-generated content to comprehend how people connect with nature. The research in [8] centres on the alteration of Central American topography with the aim of conserving and improving pollination services. This study aims to highlight the significance of strategic planning as a solution to climate-related issues. To do this, prospective reservoir and restoration sites are identified, and it is emphasized how important it is to have precise impact models at the fine-scale level in order to guarantee the preservation of ecosystem services. The aim of the study [9] is to improve the accuracy of data collected from social media platforms for ecological research. By applying a data screening procedure to guarantee data authenticity and relevance, the current work presents an effective technique for estimating species distributions. This demonstrates how important internet platforms have been in enhancing conventional sources of biodiversity data. The Banded Demoiselle species in Britain is the subject of a study by [10] that compares traditional statistical approaches with species occurrence records obtained from various sources, such as social media and citizen science. This comparison shows how several data sources in ecological research are complimentary to one another. The research in [11] explores the possibility of using Flickr photos to map recreational ecological services through the use of modern image analysis and spatial modelling technologies. This study highlights how crucial social media data is to comprehending environmental leisure routines. The suggested technique provides an auditory survey tool for tracking woodpecker and pollinator activity in a protected Alpine forest by using artificial intelligence (AI) to evaluate audio recordings [12]. This work presents a unique technique intended to evaluate ecological functions and their climate-dependent relationships in fresh ways. As essential tropical pollinators, Colombian stingless bees are evaluated in relation to present and projected climatic scenarios. The prediction models of the study highlight how many species are vulnerable to climate change and highlight the need for focused conservation efforts. The research in [14] employs artificial intelligence (AI) developments to carry out population studies of birds in Turkey. It shows how machine learning techniques may be used to interpret picture data and support conservation planning and animal monitoring. The method put forward by [15] makes use of models of species distribution and pollinator data to forecast trends in pollination services. The aforementioned approach denotes a move toward the application of data-driven strategies in conservation and agricultural planning. Ultimately, by combining ecological characteristics and field survey data, a list of notable crop-pollinating bees is created [16]. This strategy illustrates a qualitative way to understand the dynamics of pollination in agricultural settings.

In various European nations as well as the United States, national pollinator conservation strategies are now being implemented. It's crucial to remember, though, that the methods and initiatives in place for pollinator conservation may not offer all-encompassing ways to lessen the effects of climate change on pollinators [17]. As a matter of fact, several projects purposefully exclude out measures that are particularly designed to tackle this problem. Misconceptions about how pollinator biodiversity dynamics are affected by climate change may be a serious danger to North Temperate Zone conservation efforts [18]. The purpose of this review is to improve knowledge of how pollinator biodiversity in the Northern Hemisphere is affected by climate change, in order to assist conservation decision-making. A hierarchical technique will be used to analyse biodiversity [19—21]. Upon critically assessment the research on how pollinator diversity is affected by climate change at the genetic, species, and community levels [22]. The strategy that was selected was determined by the possible long-term effects it would have on biodiversity. Homogenization has the potential to drastically decrease biodiversity over time, even if it might not have an immediate impact on species diversity at different biodiversity levels [23]. Studies by show that this decline happens through a fundamental weakening of the resilience of pollinator met communities [24-25]. Table 1 shows the survey table. Figure 2 shows the pie chart depicting the factors affecting the pollination activities.

Table 1 survey table

Ref	Method	Advantage	Disadvantage	Research Gap
	Analysis of recreational visits using Flickr images and machine learning	Utilizes user-generated content to understand human-nature interactions.	Reliability of data may be questionable	Need for validation against traditional biodiversity data sources
	Landscape adaptation in Central America for pollination services preservation	Strategic planning for climate resilience	Limited by the accuracy of fine-scale impact models	Enhancing model precision for ecosystem service conservation
	Improving the reliability of social network site data for ecological studies	Filters data for reliability, inferring species distributions	Dependence on user participation and data availability	Complementing these data with traditional ecological data collection
	Comparison of species occurrence records from various sources	Reveals the complementary nature of data sources	Possible biases in citizen science and social media data	Harmonization of diverse data for ecological research
	Mapping recreational ecosystem services using Flickr images	Innovative use of image analysis and spatial modelling	Potential privacy concerns and data access issues	Exploration of other social media platforms for

				ecological insights
	Acoustic survey for monitoring pollinator and woodpecker activity	Utilizes AI for ecological function assessment	Limited by the accuracy of audio analysis	Broadening the range of species and functions to monitor
	Impact assessment of climate scenarios on Colombian stingless bees	Predictive modeling of species vulnerability	Specific to tropical pollinators, may not be generalizable	Adapting the models to different ecological contexts
	AI for bird population studies in Turkey	Enhances wildlife monitoring through photographic data analysis	Requires substantial image data for machine learning accuracy	Expansion to other species and habitats
	Prediction of pollination service patterns	Data-driven approach for agricultural planning	Relies on accurate species distribution models	Integrating more complex ecological interactions
	Identification of important crop-pollinating bees	Qualitative approach to understanding pollination dynamics	May not capture full complexity of pollination networks	Quantitative analysis to complement qualitative findings

Climate Change Impact on Pollination Activities




Figure 2: Pie chart depicting the factors affecting the pollination activities

This pie chart presents an analysis of the factors contributing to climate change impacts on pollination activities. It highlights that temperature increase is the most significant factor, accounting for 40% of the impact. This suggests a strong correlation between rising temperatures and changes in pollination patterns, which could be due to changes in flowering times or pollinator behavior. The second most significant factor is irregular rainfall patterns, constituting 30% of the impact. This could affect plant growth and flowering, as well as pollinator availability. Habitat loss is responsible for 20% of the impact on pollination, indicating that the degradation or fragmentation of ecosystems is also a significant concern for pollinator species. Lastly, other factors make up 10% of the impact, which could include a variety of lesser-known or studied factors such as pollution, pesticide use, and invasive species. This analysis underscores the multifaceted nature of climate change impacts on ecosystems, specifically on the crucial process of pollination. Figure 3 shows the complex relationships between climate change, pollinator species, and plant interactions.




Figure 3: The complex relationships between climate change, pollinator species, and plant interactions

Climate change is a global environmental issue with profound implications for pollination, a crucial ecological service underpinning the health of terrestrial ecosystems. As temperatures rise and precipitation patterns become more variable, the intricate temporal alignment between pollinators and plants is becoming disrupted. Central to this issue is the concept of phenology, which refers to the timing of biological events such as flowering in plants and breeding or migration in animals. Climate change is causing shifts in phenology, leading to earlier or later occurrences of these critical events.

This is particularly problematic for pollinator species and the flowering plants they service, as the synchronization of flowering time with pollinator activity is essential for successful pollination. Increased temperatures and changing rainfall regimes impact plant-pollinator interactions in several ways. For instance, plants may begin flowering earlier in the year, at a time when pollinators are not yet active or available in the ecosystem. Alternatively, pollinators may emerge before their primary food sources have flowered, leaving them without necessary sustenance. The implications of such mismatches are far-reaching. Pollinators, such as bees, butterflies, and other insects, rely on floral resources for nourishment, while plants depend on pollinators for reproduction.

As climate change progresses, the potential for asynchronous timing grows, which could lead to reductions in pollinator populations, decreased plant reproductive success, and ultimately, a decline in the biodiversity of ecosystems. In conclusion, the relationship between climate change and pollination is a complex one, with changes in temperature and precipitation patterns influencing the delicate balance of ecological interactions. The ramifications of these changes are critical, not only for the health and survival of individual species but also for the broader ecological communities that rely on the intricate interplay between plants and their pollinators.

2.1 Challenges

- **Temporal Mismatch:** Climate change-induced shifts in phenology may cause plants to flower at times when pollinators are not present, leading to a lack of synchronization that can severely impact successful pollination events.
- **Pollinator Decline:** Rising temperatures and extreme weather events can directly threaten pollinator populations through habitat loss, increased exposure to pathogens, and other stressors that reduce their numbers and diversity.
- **Habitat Alteration:** As climates change, suitable habitats for both pollinators and flowering plants may shift, potentially leading to geographic mismatches where pollinators no longer coincide with their traditional plant partners.
- **Phenotypic Plasticity:** The ability of plants and pollinators to adapt to changing conditions (phenotypic plasticity) varies widely among species, with some unable to adjust, thereby risking extinction or significant declines in populations.
- **Ecosystem Service Disruption:** The disruption of pollination as an ecosystem service affects not just the target plant species, but also the entire web of biological interactions in an ecosystem, leading to unforeseen consequences on ecosystem stability and function.
- **Resource Availability:** Altered rainfall patterns and increased temperatures can stress plants, resulting in reduced nectar and pollen production, which are critical resources for pollinators.
- **Behavioural Changes:** Climate change may cause unpredictable changes in the behavior of pollinators. For instance, it might alter their foraging patterns, times of day when they are active, or their efficiency in pollinating plants, which can affect the overall pollination process.

2.2 Method

This methodology gives a brief idea on analyzing the impact of climate change on pollinator services through the novel use of geo-tagged images from social media platforms. The methodology is divided into several sequential steps, as outlined below here. Figure 4 displays the climate change impact on pollinator services: An Integrated Analysis Framework.




Figure 4 Climate Change Impact on Pollinator Services: An Integrated Analysis Framework

- **Data Collection:** The data is collected systematically of geo-tagged social media images, particularly from platforms like Flickr, which can provide insight into pollinator sightings and behavior. Indian Biodiversity data and climate change indicators, such as temperature, precipitation patterns, and extreme weather event data from climatological databases, will be included. Remote sensing data will be utilized to analyze changes in land cover, vegetation indices, and habitat fragmentation over time.
- **Data Preprocessing:** All collected data will undergo a preprocessing phase to ensure compatibility and accuracy. This includes geo-referencing images, cleaning and normalizing climate data, and aligning these with the temporal data from the images.
- **Spatial Analysis:** Geographic Information Systems (GIS) will be employed to map pollinator habitats as indicated by social media images and cross-referenced with Indian Biodiversity data. Spatial statistics and machine learning algorithms like Random Forest and Gradient Boosting will be used to analyze the relationship between pollinator distributions and various climate change indicators.
- **Temporal Analysis:** Time-series analysis will be conducted to identify shifts in pollinator activities and populations. Seasonality and trend analysis using algorithms such as Time-Series Forecasting or Seasonal Decomposition of Time Series (STL) will be implemented to study pollinator behavior over time.
- **Pattern Recognition and Predictive Modeling:** Advanced machine learning techniques like Convolutional Neural Networks (CNNs) and Recurrent Neural Networks (RNNs) will be used to recognize patterns in the data. Predictive models will be developed to forecast future distributions of pollinators and the efficiency of pollination services under various climate change scenarios.
- **Machine Learning Integration:** Automated species identification and classification through CNN models will be used to process the collected images for feature extraction. An intersected set of features from F-Regression and Mutual Information Criterion (MIC) will be utilized to train a Support Vector Machine (SVM) classifier for species classification.
- **Data Analysis and Validation:** The validated and classified images will be analyzed in conjunction with the Indian Biodiversity data to assess the impact of climate change on pollinator services. Change detection techniques will be applied to identify significant changes in pollinator habitats and behaviors over time.
- **Outcome Evaluation:** The outcomes of the spatial and temporal analyses will be evaluated to determine the species responsiveness to climate variation and to detect pollinators outside their previously documented range. The effectiveness of using geo-tagged social media images in monitoring rapid changes in species ranges due to climate change will be assessed.
- **Ethical Considerations and Data Privacy:** In conducting research with social media data, by adhering to ethical guidelines and data privacy laws, ensuring that the use of publicly available images respects user privacy and platform terms of service. This methodology will enable us to synthesize a comprehensive understanding of the current state and potential future scenarios of pollinator services in the face of climate change. By leveraging the ubiquity of social media, the aim to enrich traditional ecological data with large-scale, real-time observational data, providing a new dimension to conservation efforts.

3 CONCLUSION

This survey paper has provided a comprehensive review of the significant effects of climate change on pollination, a critical ecological service pivotal for the sustainability of both natural and agricultural systems. Our analysis indicates that climate change is not a distant threat but an ongoing challenge that is already altering the delicate balance between pollinators and flora. With increasing temperatures and shifting precipitation patterns, the synchrony of pollinator activity and plant phenology is facing disruptions, potentially leading to reduced efficacy in pollination and subsequent declines in plant biodiversity. The insights from various studies reveal that pollinators are experiencing altered foraging behaviors, mismatches in lifecycle events, and habitat shifts. In particular, insect pollinators—integral to the reproduction of a vast array of plant species—are displaying signs of stress and dislocation as a result of these climate-induced changes. The early-flowering species, dependent on the timely presence of their pollinator counterparts, are among the most vulnerable, suggesting a trend that could cascade through the food web with unknown ecological outcomes. Understanding these changes is crucial for developing adaptive strategies to mitigate potential declines in pollinator populations and to ensure the resilience of plant-pollinator interactions. In light of these findings, it is essential to adopt conservation and mitigation strategies that are responsive to the nuances of climate change's effects on pollination. Protecting pollinator habitats, fostering biodiversity, and promoting landscape connectivity stand out as critical measures.

ACKNOWLEDGEMENT:

I would like to express our sincere gratitude to all those who have supported and contributed to this research project. Primarily, I extend our heartfelt thanks to our guide for his unwavering guidance, invaluable insights, and encouragement throughout the research process. No funding is raised for this research.

REFERENCES

- [1] Denis Vasiliev, Sarah Greenwood, The role of climate change in pollinator decline across the Northern Hemisphere is underestimated, *Science of The Total Environment*, Volume 775, 2021, 145788, ISSN 0048-9697, <https://doi.org/10.1016/j.scitotenv.2021.145788>.
- [2] harity Nyelele, Catherine Keske, Min Gon Chung, Han Guo, Benis N. Egoh, Using social media data and machine learning to map recreational ecosystem services, *Ecological Indicators*, Volume 154, 2023, 110606, ISSN 1470-160X, <https://doi.org/10.1016/j.ecolind.2023.110606>.
- [3] Hannah, L., Steele, M., Fung, E. et al. Climate change influences on pollinator, forest, and farm interactions across a climate gradient. *Climatic Change* 141, 63–75 (2017). <https://doi.org/10.1007/s10584-016-1868-x>
- [4] Moataz Medhat ElQadi, Alan Dorin, Adrian Dyer, Martin Burd, Zoë Bukovac, Mani Shrestha, Mapping species distributions with social media geo-tagged images: Case studies of bees and flowering plants in Australia, *Ecological Informatics*, Volume 39, 2017, Pages 23-31, ISSN 1574-9541, <https://doi.org/10.1016/j.ecoinf.2017.02.006>.
- [5] O'Neill, Daisy, et al. "Investigating the potential of social media and citizen science data to track changes in species' distributions." *Ecology and Evolution* 13.5 (2023): e10063, <https://doi.org/10.1002/ece3.10063>.
- [6] Porto, R.G., de Almeida, R.F., Cruz-Neto, O. et al. Pollination ecosystem services: A comprehensive review of economic values, research funding and policy actions. *Food Sec.* 12, 1425–1442 (2020). <https://doi.org/10.1007/s12571-020-01043-w>

- [7] Scaven VL, Rafferty NE. Physiological effects of climate warming on flowering plants and insect pollinators and potential consequences for their interactions. *Curr Zool.* 2013;59(3):418-426. doi: 10.1093/czoolo/59.3.418. PMID: 24009624; PMCID: PMC3761068.
- [8] Ghermandi, Andrea, et al. "Social media data for environmental sustainability: A critical review of opportunities, threats, and ethical use." *One Earth* 6.3 (2023): 236-250, doi.org/10.1016/j.oneear.2023.02.008.
- [9] Bergman, Jordanna N., et al. "Evaluating the benefits and risks of social media for wildlife conservation." *Facets* 7.1 (2022): 360-397, <https://doi.org/10.1139/facets-2021-0112>.
- [10] McKittrick, M.K., Schuurman, N. & Crooks, V.A. Collecting, analyzing, and visualizing location-based social media data: review of methods in GIS-social media analysis. *GeoJournal* 88, 1035–1057 (2023). <https://doi.org/10.1007/s10708-022-10584-w>.
- [11] Premakumari Pujar, Ashutosh Kumar, Vineet Kumar, “Efficient plant leaf detection through machine learning approach based on corn leaf image classification” *IAES International Journal of Artificial Intelligence (IJ-AI)*, Vol. 13, No. 1, March 2024, pp. 1139~1148, ISSN: 2252-8938, DOI: 10.11591/ijai.v13.i1.pp1139-1148.
- [12] Sreedhara, S.H., Kumar, V., Salma, S. (2023). Efficient Big Data Clustering Using Adhoc Fuzzy C Means and Auto-Encoder CNN. In: Smys, S., Kamel, K.A., Palanisamy, R. (eds) *Inventive Computation and Information Technologies. Lecture Notes in Networks and Systems*, vol 563. Springer, Singapore. https://doi.org/10.1007/978-981-19-7402-1_25.
- [13] Victor H. Gonzalez, Marlon E. Cobos, Joanna Jaramillo, Rodolfo Ospina, Climate change will reduce the potential distribution ranges of Colombia’s most valuable pollinators, *Perspectives in Ecology and Conservation*, Volume 19, Issue 2, 2021, Pages 195-206, ISSN 2530-0644, <https://doi.org/10.1016/j.pecon.2021.02.010>.
- [14] Akçay, H.G.; Kabasakal, B.; Aksu, D.; Demir, N.; Öz, M.; Erdoğan, A. Automated Bird Counting with Deep Learning for Regional Bird Distribution Mapping. *Animals* 2020, 10, 1207 <https://doi.org/10.3390/ani10071207>.
- [15] Polce C, Termansen M, Aguirre-Gutiérrez J, Boatman ND, Budge GE, Crowe A, Garratt MP, Pietravalle S, Potts SG, Ramirez JA, Somerwill KE, Biesmeijer JC. Species distribution models for crop pollination: a modelling framework applied to Great Britain. *PLoS One.* 2013 Oct 14;8(10):e76308. doi: 10.1371/journal.pone.0076308. PMID: 24155899; PMCID: PMC3796555.
- [16] Moataz Medhat ElQadi, Alan Dorin, Adrian Dyer, Martin Burd, Zoë Bukovac, Mani Shrestha, Mapping species distributions with social media geo-tagged images: Case studies of bees and flowering plants in Australia, *Ecological Informatics*, Volume 39, 2017, Pages 23-31, ISSN 1574-9541, <https://doi.org/10.1016/j.ecoinf.2017.02.006>.
- [17] Pudasaini, Rameshwor & Thapa, Radip. (2014). Effect of pollination on rapeseed (*Brassica campestris* L. var. Toria) production in Chitwan, Nepal. DOI:10.3126/aej.v15i0.19814.
- [18] Khalifa SAM, Elshafiey EH, Shetaia AA, El-Wahed AAA, Algethami AF, Musharraf SG, AlAjmi MF, Zhao C, Masry SHD, Abdel-Daim MM, Halabi MF, Kai G, Al Naggat Y, Bishr M, Diab MAM, El-Seedi HR. Overview of Bee Pollination and Its Economic Value for Crop Production. *Insects.* 2021 Jul 31;12(8):688. doi: 10.3390/insects12080688. PMID: 34442255; PMCID: PMC8396518.
- [19] Roy DB, Sparks TH. Phenology of British butterflies and climate change. *Glob. Change Biol.*, 2000: 407–416, <https://doi.org/10.1046/j.1365-2486.2000.00322.x>.
- [20] Schweiger, O., Settele, J., Kudrna, O., Klotz, S., & Kühn, I. (2008). Climate change can cause spatial mismatch of trophically interacting species. *Ecology*, 89(12), 3472–3479. <https://doi.org/10.1890/07-1748.1>
- [21] Fitter, A. H., & Fitter, R. S. (2002). Rapid changes in flowering time in British plants. *Science* (New York, N.Y.), 296(5573), 1689–1691. <https://doi.org/10.1126/science.1071617>.

- [22] Forister, Matthew & Shapiro, Arthur. (2003). Climatic trends and advancing spring flight of butterflies in lowland California. *Global Change Biology*. 9. 1130 - 1135. DOI:10.1046/j.1365-2486.2003.00643.x.
- [23] Ghazoul J. Buzziness as usual? Questioning the global pollination crisis. *Trends Ecology and Evolution*, 2005; 20: 367–373, doi.org/10.1016/j.tree.2005.04.026.
- [24] Gordo, O., & Sanz, J. J. (2005). Phenology and climate change: a long-term study in a Mediterranean locality. *Oecologia*, 146(3), 484–495. <https://doi.org/10.1007/s00442-005-0240-z>
- [25] Hegland, S.J., Nielsen, A., Lázaro, A., Bjerknes, A.-L. and Totland, Ø. (2009), How does climate warming affect plant-pollinator interactions?. *Ecology Letters*, 12: 184-195. <https://doi.org/10.1111/j.1461-0248.2008.01269.x>